



WEST BENGAL STATE UNIVERSITY
B.Sc. Honours 5th Semester Examination, 2022-23

PHSACOR11T-PHYSICS (CC11)

QUANTUM MECHANICS AND APPLICATIONS

Full Marks: 40

Time Allotted: 2 Hours

*The figures in the margin indicate full marks.
Candidates should answer in their own words and adhere to the word limit as practicable.
All symbols are of usual significance.*

Question No. 1 is compulsory and answer any two from the rest

1. Answer any *ten* questions from the following: 2×10 = 20
- Explain the physical significance of energy time uncertainty relation.
 - Give the physical interpretation of the wave function $\psi(x, t)$.
 - What are stationary states in quantum mechanics?
 - If the commutation relation between x and p is $[x, p] = i\hbar$. Find the commutation value of $[x^2, p]$.
 - What is the implication of the result: $[\hat{H}, \hat{L}] = 0$?
 - Consider the operator $\hat{Q} = i \frac{d}{d\phi}$ where ϕ is usual polar coordinates in two dimensions. Write down its eigenvalue equation and find its eigenvalues.
 - What is normal Zeeman effect? Under what conditions it may be observed?
 - What is Larmor precession of electron in an atom?
 - Explain bound and unbound states in quantum mechanics.
 - A wavefunction ψ is constructed as a linear combination of a set of orthonormal eigenfunctions ψ_n :

$$\psi = \sum_{n=1}^{\infty} c_n \psi_n$$

where c_n are constants. Show that if ψ is normalized then $\sum_{n=1}^{\infty} |c_n|^2 = 1$

- If the wavefunction of a particle trapped in space between $x=0$ and $x=L$ is given by $\psi(x) = A \sin \frac{2\pi x}{L}$, where A is a constant, for which value(s) of x will the probability of finding the particle be maximum?
- Electron configuration of Sodium is given by $1s^2 2s^2 2p^6 3s^1$. Find the ground state term symbol of Sodium.
- What is Stark effect?

- (n) A beam of spin $\frac{1}{2}$ particle is prepared in the state $|\psi\rangle = \frac{3}{\sqrt{34}}|+\rangle + i\frac{5}{\sqrt{34}}|-\rangle$;
where $|+\rangle$ and $|-\rangle$ are eigen states of \hat{S}_z with eigenvalues $+\frac{\hbar}{2}$ and $-\frac{\hbar}{2}$
respectively. Find the average value in S_z measurement.

2. (a) If ψ_1 and ψ_2 are two eigen states with energy E_1 and E_2 respectively, check whether the state $(\psi_1 + \psi_2)$ is stationary or not. 2

- (b) (i) Prove that the time rate of change of the expectation value of a dynamical variable satisfies the following relation 3+2

$$\frac{d}{dt}\langle\hat{A}\rangle = -\frac{i}{\hbar}\langle[\hat{A}, \hat{H}]\rangle + \left\langle\frac{\partial\hat{A}}{\partial t}\right\rangle$$

where the symbols have their usual meanings.

- (ii) Using the above relation show that the time rate of change of expectation value of momentum is equal to the average value of force.

- (c) Prove that the parity of spherical harmonics $Y_{l,m}(\theta, \phi)$ is $(-1)^l$. 2

- (d) What do you mean by degenerate wavefunction? 1

3. (a) The potential in a region is given as:

$$\begin{aligned} V(x) &= 0 \text{ for } x < 0 \\ &= V_0 \text{ for } 0 \leq x \leq a \\ &= 0 \text{ for } x > a \end{aligned}$$

A particle of mass m and energy $E < V_0$ travelling from left to the right is incident on the potential barrier.

- (i) Write down Schrodinger equations in three regions of the potential. 2

- (ii) Write down appropriate boundary conditions. 2

- (b) The wavefunction of a hydrogen atom is given by the following superposition of energy eigenfunctions $\psi_{nlm}(\vec{r})$ (n, l, m are the usual quantum numbers):

$$\psi(\vec{r}) = \sqrt{\frac{2}{7}}\psi_{100}(\vec{r}) - \frac{3}{\sqrt{7}}\psi_{210}(\vec{r}) + \frac{1}{\sqrt{14}}\psi_{322}(\vec{r})$$

- (i) Determine the ratio of expectation value of the energy to the ground state energy. 2

- (ii) What are the expectation value of \hat{L}^2 and \hat{L}_z operators? 2

- (iii) What is the probability that the atom is found in a state of even parity? 2

4. (a) Hamiltonian for the linear harmonic oscillator is given by $\hat{H} = \frac{1}{2}\hat{p}^2 + \frac{1}{2}m\omega^2\hat{x}^2$,
where the symbols have usual meanings. Using the basic commutation relation between \hat{x} and \hat{p} show that,

- (i) $[\hat{a}, \hat{a}^+] = 1$ and 2+2

(ii) the Hamiltonian is given by

$$H = (\hat{a}^+ \hat{a} + 1/2) \hbar \omega$$

$$\text{given that } \hat{a} = \left(\frac{m\omega}{2\hbar} \right)^{1/2} \left(\hat{x} + \frac{i}{m\omega} \hat{p} \right) \text{ and } \hat{a}^+ = \left(\frac{m\omega}{2\hbar} \right)^{1/2} \left(\hat{x} - \frac{i}{m\omega} \hat{p} \right)$$

Then find the normalized ground state wavefunction of linear harmonic oscillator.

(b) A particle constrained to move along x -axis in the domain $0 \leq x \leq L$ has a wavefunction $\psi(x) = \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi x}{L}\right)$, where n is an integer. What is the expectation value of its momentum?

5. (a) State Moseley's Law. Derive this law from Bohr's theory.

(b) Considering the L-S coupling scheme for helium atom, find the spectroscopic terms for (i) $1s^1 2s^1$ and (ii) $1s^1 2p^1$ configurations.

(c) In a Stern-Gerlach experiment on turning on the magnetic field, the beam splits into seven components. What is the angular momentum of the atoms in the beam?

—x—